

# Lattices and the Knaster-Tarski Theorem

Deepak D'Souza


Department of Computer Science and Automation  
Indian Institute of Science, Bangalore.

8 August 2018


# Outline

- 1 Why study lattices
- 2 Partial Orders
- 3 Lattices
- 4 Knaster-Tarski Theorem
- 5 Computing LFP

## What a lattice looks like



Subsets of  $\{1, 2, 3\}$ ,  
"subset"



Odd/even, "contained  
in"

## Why study lattices in program analysis?

### Why lattices?

- Natural way to obtain the “collecting state” at a point is to take union of states reached along each path leading to the point.
- With abstract states also we want a “union” or “join” over all paths (JOP).


```

1: p := 5;
2: q := 2;
3: while (p > q) {
4:   p := p+1;
5:   q := q+2;
6: }
6: print p;

```


### Why fixpoints?

- Guaranteed to safely approximate JOP (\* Conditions apply).
- Easier to compute than JOP.
- Knaster-Tarski theorem tells us when a fixpoint exists and what it looks like.




## Partial Orders


- Usual order (or **total** order) on numbers:  $1 \leq 2 \leq 3$ .
- Some domains are naturally “partially” ordered:



Subsets of  $\{1, 2, 3\}$ ,  
“subset”



Divisors of 12, “divides”



Odd/even, “contained  
in”

## Partial orders: definition


- A **partially ordered set** is a non-empty set  $D$  along with a partial order  $\leq$  on  $D$ . Thus  $\leq$  is a binary relation on  $D$  satisfying:
  - $\leq$  is reflexive ( $d \leq d$  for each  $d \in D$ )
  - $\leq$  is transitive ( $d \leq d'$  and  $d' \leq d''$  implies  $d \leq d''$ )
  - $\leq$  is anti-symmetric ( $d \leq d'$  and  $d' \leq d$  implies  $d = d'$ ).

## Binary relations as Graphs

We can view a binary relation on a set as a **directed graph**.  
For example, the binary relation

$$\{(a, a), (a, b), (b, c), (b, e), (d, e), (d, c), (e, f)\}$$


can be represented as the graph:




## Partial Order as a graph

A **partial order** is then a special kind of directed graph:

- Reflexive = self-loop on each node
- Antisymmetric = no 2-length cycles
- Transitive = “transitivity” of edges.



Graph  
representation




Hasse-diagram  
representation



## Upper bounds etc.




In a partially ordered set  $(D, \leq)$ :

- An element  $u \in D$  is an **upper bound** of a set of elements  $X \subseteq D$ , if  $x \leq u$  for all  $x \in X$ .
- $u$  is the **least upper bound** (or **lub** or **join**) of  $X$  if  $u$  is an upper bound for  $X$ , and for every upper bound  $y$  of  $X$ , we have  $u \leq y$ . We write  $u = \bigsqcup X$ .
- Similarly,  $v = \bigsqcap X$  ( $v$  is the **greatest lower bound** or **glb** or **meet** of  $X$ ).






# Lattices

- A **lattice** is a partially order set in which every pair of elements has an lub and a glb.
- A **complete** lattice is a lattice in which every **subset** of elements has a lub and a glb.




# Lattices

- A **lattice** is a partially order set in which every pair of elements has an lub and a glb.
- A **complete** lattice is a lattice in which every **subset** of elements has a lub and a glb.
- Examples below are all complete lattices.



## More lattices




## Exercise

- 1 Example of a partial order that is not a lattice?


## Exercise

- ① Example of a partial order that is not a lattice?



## Exercise


- 1 Example of a partial order that is not a lattice?



- 2 “Simplest” example of a partial order that is not a lattice?

## Exercise

- 1 Example of a partial order that is not a lattice?




- 2 “Simplest” example of a partial order that is not a lattice?





## Exercise

- 1 Example of a partial order that is not a lattice?




- 2 “Simplest” example of a partial order that is not a lattice?




- 3 Example of a lattice which is **not** complete?

## Exercise

- 1 Example of a partial order that is not a lattice?



- 2 “Simplest” example of a partial order that is not a lattice?




- 3 Example of a lattice which is **not** complete?



## Monotonic functions

Let  $(D, \leq)$  be a partially ordered set.



- A function  $f : D \rightarrow D$  is **monotonic** or **order-preserving** if whenever  $x \leq y$  we have  $f(x) \leq f(y)$ .



## Partial order induced by a subset of elements


Let  $(D, \leq)$  be a partially ordered set, and  $X$  be a non-empty subset of  $D$ . Then  $X$  induces a partial order, which we call the partial order *induced by  $X$*  in  $(D, \leq)$ , and defined to be  $(X, \leq \cap (X \times X))$ .

Example: the partial order induced by the set of elements  $X = \{2, 3, 12\}$ .



## Fixpoints

- A **fixpoint** of a function  $f : D \rightarrow D$  is an element  $x \in D$  such that  $f(x) = x$ .
- A **pre-fixpoint** of  $f$  is an element  $x$  such that  $x \leq f(x)$ .
- A **post-fixpoint** of  $f$  is an element  $x$  such that  $f(x) \leq x$ .




## Knaster-Tarski Fixpoint Theorem

### Theorem (Knaster-Tarski)

Let  $(D, \leq)$  be a complete lattice, and  $f : D \rightarrow D$  a monotonic function on  $(D, \leq)$ . Then:

- (a)  $f$  has at least one fixpoint.
- (b)  $f$  has a **least fixpoint** which coincides with the glb of the set of postfixpoints of  $f$ , and a **greatest fixpoint** which coincides with the lub of the prefixpoints of  $f$ .
- (c) The set of fixpoints  $P$  of  $f$  itself forms a complete lattice under  $\leq$ .

## Fixpoints of $f$




Stars denote fixpoints.

## Exercise

Consider the complete lattice and monotone function  $f$  below.

- 1 Mark the pre-fixpoints with up-triangles ( $\triangle$ ).
- 2 What is the lub of the pre-fixpoints?
- 3 Mark post-fixpoints with down-triangles ( $\nabla$ ).
- 4 Fixpoints are the stars ( $\boxtimes$ ).




Check that claims of K-T theorem hold here.



## Exercise

Consider the complete lattice and monotone function  $f$  below.

- 1 Mark the pre-fixpoints with up-triangles ( $\triangle$ ).
- 2 What is the lub of the pre-fixpoints?
- 3 Mark post-fixpoints with down-triangles ( $\nabla$ ).
- 4 Fixpoints are the stars ( $\boxtimes$ ).




Check that claims of K-T theorem hold here.

If you drop completeness of lattice or monotonicity of  $f$ , does K-T still hold?

## Proof of Knaster-Tarski theorem


- (a)  $g = \bigsqcup Pre$  is a fixpoint of  $f$ .
- (b)  $g$  is the greatest fixpoint of  $f$ .
- (c) Similarly  $l = \bigsqcap Post$  is the least fixpoint of  $f$ .
- (d) Let  $P$  be the set of fixpoints of  $f$ . Then  $(P, \leq)$  is a *complete* lattice.

## Proof of K-T theorem: (a)



## Proof of K-T theorem: (a)


To show  $g = f(g)$ :



## Proof of K-T theorem: (a)

To show  $g = f(g)$ :


- $g \leq f(g)$



## Proof of K-T theorem: (a)

To show  $g = f(g)$ :


- $g \leq f(g)$ 
  - Since  $f(g)$  can be seen to be u.b. of  $Pre$ .



## Proof of K-T theorem: (a)

To show  $g = f(g)$ :


- $g \leq f(g)$ 
  - Since  $f(g)$  can be seen to be u.b. of  $Pre$ .
- $f(g) \leq g$



## Proof of K-T theorem: (a)

To show  $g = f(g)$ :


- $g \leq f(g)$ 
  - Since  $f(g)$  can be seen to be u.b. of  $Pre$ .
- $f(g) \leq g$ 
  - Since  $f(g)$  can be seen to be prefixpoint of  $f$ .






## Proof of K-T theorem: (b)

$g$  is the greatest fixpoint of  $f$ .



## Proof of K-T theorem: (b)

$g$  is the greatest fixpoint of  $f$ .  
Any other fixpoint is also a  
pre-fixpoint of  $f$ , and hence  $g$   
must dominate it.




## Exercise: intervals and closure

Let  $(D, \leq)$  be a partial order, and let  $f : D \rightarrow D$ .

- Let  $a, b \in D$ . The **interval** from  $a$  to  $b$ , written  $[a, b]$ , is the set  $\{d \mid a \leq d \leq b\}$ .
- A subset  $X \subseteq D$  is said to be **closed** wrt to  $f$ , if  $f(x) \in X$  for each  $x \in X$ .


Exercise: Let  $(D, \leq)$  be a partial order with a  $\top$  element, and let  $f : D \rightarrow D$  be a monotone function on  $D$ .

- 1 Show that an interval in  $D$  need *not* be closed wrt  $f$ .
- 2 Let  $u \in D$  be the lub of a set  $X$  of fixpoints of  $f$ . Prove that the interval  $[u, \top]$  is closed wrt  $f$ .



## Proof of K-T theorem: (d)

- $(P, \leq)$  is also a partial order.
- $(P, \leq)$  is a complete lattice
  - Let  $X \subseteq P$ . We show there is an lub of  $X$  in  $(P, \leq)$ .
    - Let  $u$  be lub of  $X$  in  $(D, \leq)$ .
    - Consider “interval”  $I = [u, \top] = \{x \in D \mid u \leq x\}$ .  $(I, \leq)$  is also a complete lattice.
    - $f : I \rightarrow I$  as well, and monotonic on  $(I, \leq)$ .
    - Hence by part (a)  $f$  has a least fixpoint in  $I$ , say  $v$ .
    - Argue that  $v$  is the lub of  $X$  in  $(P, \leq)$ .



## Chains in partial orders

- A **chain** in a partial order  $(D, \leq)$  is a totally ordered subset of  $D$ .
- An **ascending chain** is an infinite sequence of elements of  $D$  of the form:

$$d_0 \leq d_1 \leq d_2 \leq \dots .$$

- An ascending chain  $\langle d_i \rangle$  is **eventually stable** if there exists  $n_0$  such that  $d_i = d_{n_0}$  for each  $i \geq n_0$ .
- $(D, \leq)$  has **finite** height if each chain in it is finite.
- $(D, \leq)$  has **bounded** height if there exists  $k$  such that each chain in  $D$  has height at most  $k$  (i.e. number of elements in each chain is at most  $k + 1$ .)

## Monotonicity, distributivity, and continuity

- $f$  is monotone:

$$x \leq y \implies f(x) \leq f(y).$$

- $f$  is distributive:


$$f(x \sqcup y) = f(x) \sqcup f(y).$$

- $f$  is continuous: For any asc chain  $X$ :


$$f(\bigsqcup X) = \bigsqcup (f(X)).$$

- $f$  is inf distributive: For any  $X \subseteq D$ :


$$f(\bigsqcup X) = \bigsqcup (f(X)).$$




Monotonic



Distributive



Continuous



Inf-Distributive

# Characterising lfp's and gfp's of a function $f$ in a complete lattice $(D, \leq)$

- $f$  is **continuous** if for any ascending chain  $X$  in  $D$ ,


$$f(\bigsqcup X) = \bigsqcup (f(X)).$$

- If  $f$  is **continuous** then

$$lfp(f) = \bigsqcup_{i \geq 0} (f^i(\perp)).$$


- If  $f$  is **monotonic** and  $(D, \leq)$  has **finite height** then we can compute  $lfp(f)$  by finding the stable value of the asc. chain

$$\perp \leq f(\perp) \leq f^2(\perp) \leq f^3(\perp) \leq \dots$$



## Exercise

Consider the statement “ $p := p + q$ ”. Show the transfer function of this statement in the parity lattice below.



Is it monotonic/distributive/continuous?